

GOES-18 SEISS EHS Level 1b (L1b) Data Release  
Provisional Data Quality  
November 15, 2022  
Read-Me for Data Users

The Peer Stakeholder - Product Validation Review (PS-PVR) for the GOES-18 Space Environment In-Situ Suite (SEISS) Energetic Heavy Ion Sensor (EHIS) L1b Provisional Maturity was held on November 15, 2022. As a result of this review, the PS-PVR panel chair declared that the GOES-18 EHS L1b data are at Provisional Validation Maturity as of November 15, 2022.

The L1b data product consists of 5-minute-cadence differential directional fluxes and associated systematic (instrumental) and statistical errors. Fluxes are produced for hydrogen (H) and helium (He); for the carbon-nitrogen-oxygen (CNO), neon-sulfur (Ne-S), and chlorine-nickel (Cl-Ni) mass groups; and for individual elements between beryllium and copper (Be-Cu) (but see below for a restriction on the beryllium and boron fluxes). EHS has a single 60 degree (full cone angle) field-of-view directed radially outward from the Earth (toward zenith). The energy range is nominally 10-200 MeV/nucleon for hydrogen (protons) and helium (alpha particles), divided into five energy channels. The energy range increases with atomic number (Z) since the stopping power in silicon is the same for all species in each energy channel. Outside of solar energetic particle (SEP) events, EHS observes backgrounds, some of which may be from galactic cosmic ray (GCR) fluxes.

The H and He fluxes are derived directly from coincidence rates (3-second cadence in the raw Level 0 data), as with SGPS, and can be averaged over longer periods to improve the counting statistics. However, the heavy ion fluxes are derived using a maximum likelihood (ML) fit to a histogram of Z values determined on-orbit (sum of five 1-minute cadence histograms) using the angle-detecting inclined sensor (ADIS) system incorporated into the EHS telescope. While this ML fit is necessary for meeting requirements in the presence of very sparse heavy ion count rates, it limits the utility of the L1b data in post-processing. Moreover, when the lower one-sigma statistical error is equal to the mean value, only an upper limit exists (mean plus upper one-sigma statistical error). (In the L1b files, the mean fluxes are contained in the variable 'BeCu5MinuteDifferentialFluxes', and the lower and upper statistical errors are contained in the variable 'BeCu5MinuteDifferentialFluxStatErrorsBounds'.) Averaging ML fits will not improve the statistics and is invalid when the set of fits includes upper limits. As a result, derivation of heavy ion fluxes for periods longer than 5 minutes requires reprocessing from Level 0 raw data. This is a limitation that is independent of the maturity of the product.

Provisional validation means:

- Validation activities are ongoing and the general research community is now encouraged to participate.
- Severe algorithm anomalies are identified and under analysis. Solutions to anomalies are in development and testing.
- Incremental product improvements may still be occurring.
- Product performance has been demonstrated through analysis of a small number of independent measurements by another instrument on the same observatory, or by other observatories.
- Product analysis is sufficient to establish product performance relative to expectations (Performance Baseline).
- Documentation of product performance exists that includes recommended remediation

strategies for all anomalies and weaknesses. Any algorithm changes associated with severe anomalies have been documented, implemented, and tested.

- Testing has been fully documented.
- Product is ready for operational use and for use in comprehensive calibration/validation activities and product optimization.

Users of the GOES-18 EHS L1b data bear responsibility for inspecting the data and understanding the known caveats prior to use. Below is the list of caveats that have been identified and are under analysis. Solutions are in development and testing.

1. Current understanding of GOES-18 EHS H and He performance is based on analysis of three weak SEP events that occurred between April and September 2022, during which significant heavy ion fluxes were not observed.
2. Current understanding of GOES-18 EHS heavy ion performance is based on analysis of histograms accumulated from July 22 to October 28, 2022. Based on this analysis, the histograms for channels E4 and E5 are inconsistent with GOES-16 and GOES-17. More on-orbit data are needed to update the tables and thereby resolve this inconsistency.
3. No EHS L1b data processed prior to declaration of Provisional Maturity (e.g., those available from CLASS) should be used, owing to important updates to flight science tables and the ground LUT for Provisional Maturity. The first data processed with consistent Provisional-maturity flight science tables and ground look-up tables (LUTs) were at 2104 UTC on November 16, 2022.
4. Absolute calibrations are still being refined. For example, a temperature sensitivity that may impart a significant diurnal variation onto the observed proton (hydrogen) fluxes remains to be quantified.
5. EHS hydrogen and helium fluxes are low relative to SGPS. The root cause of this discrepancy has not been determined. In general, NOAA recommends that the SGPS proton (hydrogen) and alpha particle (helium) fluxes be used instead of the EHS helium channels. SGPS produces hydrogen and helium fluxes over a wider energy range and in more energy channels than EHS. However, since EHS has an order-of-magnitude greater event-to-background ratio than SGPS, studies of weak events should include EHS data, keeping in mind the absolute calibration differences.
6. Lithium, beryllium, and boron ions of solar origin are never observed, being destroyed in the solar interior. Therefore, L1b Be and B fluxes are replaced with fill values because they should never be used. (Lithium fluxes are not within the reported measurement range.) It is believed that counts at these histogram locations are due to slow helium nuclei being counted as heavy ions.
7. Outside of SEP events, EHS observes backgrounds, some of which may be from GCR fluxes. Under these conditions, the L1b fluxes are not accurate, since the processing uses geometrical factors and energy bandwidths derived for SEP spectra. Steep SEP spectra emphasize the lower-energy ends of the energy response functions while flatter GCR spectra more strongly emphasize the higher-energy tails of the response functions.
8. As described above, time-averaging L1b heavy ion fluxes, particularly those that are upper limits, does not result in improved accuracy and therefore should not be performed.

Contact for further information: OSPO User Services at [SPSD.UserServices@noaa.gov](mailto:SPSD.UserServices@noaa.gov)

NCEI contacts for specific information on the EHS L1b data:

Juan Rodriguez [juan.rodriquez@noaa.gov](mailto:juan.rodriquez@noaa.gov)

Brian Kress [brian.kress@noaa.gov](mailto:brian.kress@noaa.gov)

Pamela Wyatt [pamela.wyatt@noaa.gov](mailto:pamela.wyatt@noaa.gov)

NCEI website for GOES-R Space Weather data (provides daily aggregations of EHS L1b data):

<https://www.ngdc.noaa.gov/stp/satellite/goes-r.html>